



## 日本入国時に必要な検査証明書の要件（検体、検査方法、検査時間）

※ 有効な検体、検査方法等が記載された検査証明書のみ有効と取り扱います。

<有効な検査証明書として認められる要件>

<有効な検査証明書として認められない主な例>

### 検体

- Nasopharyngeal/Nasopharynx (Swab/smear)
- Rhinopharyngeal/Rhinopharynx (Swab/Smear)**  
(鼻咽頭ぬぐい)
- (Deep throat) Saliva (唾液)
- **Nasopharyngeal (※) and oropharyngeal(throat) (swabs /smear)**  
(鼻咽頭ぬぐい・咽頭ぬぐいの混合)  
(Naso and oropharyngeal/Rhino and oropharyngeal/oro and nashopharyngeal (※))  
(※Nasopharyngeal/Rhinopharyngeal/Nasopharynx/Rhinopharynx)

- × Nasal (swab/smear) (鼻腔ぬぐい)
- × Oral (swab/smear) (口腔ぬぐい)
- × Throat (swab/smear) (咽頭ぬぐい)
- × Mid-Turbinate Nasal (swab/smear) (中鼻甲介ぬぐい)
- × Nose (鼻ぬぐい)
- × Gargle Water (うがい液)
- × mixture of sample "A" and "B"  
(「鼻咽頭ぬぐいと咽頭ぬぐいの混合検体」を除く、複数箇所から採取した検体の混合検体)  
(なお、“A”、“B”は検体を指す)

<Example>

- × Nasal and throat (swab/smear) (鼻腔・咽頭ぬぐい)
- × Pharyngeal and nasal (swab/smear) (咽頭・鼻腔ぬぐい)
- × Nasal and oropharyngeal/oropharynx (swab/smear)  
(鼻腔・口腔咽頭ぬぐい)

### 検査方法

- ※下記の方法、もしくは検査手法名を含む方法
- Nucleic acid amplification test ( (RT-) PCR)  
核酸増幅検査 ( (RT-) PCR法)  
(real-time (RT-)PCR、Q-PCR、Fluorescence-PCR、Multiplex-PCR)
  - Nucleic acid amplification test (LAMP)  
核酸増幅検査 (LAMP法) (RT-LAMP)
  - Nucleic acid amplification test (TMA)  
核酸増幅検査 (TMA法)
  - Nucleic acid amplification test (TRC)  
核酸増幅検査 (TRC法)
  - Nucleic acid amplification test (Smart Amp)  
核酸増幅検査 (Smart Amp法)
  - Nucleic acid amplification test (NEAR)  
核酸増幅検査 (NEAR法) (例: ID-NOW®)
  - Next generation sequence  
次世代シーケンス法
  - Quantitative antigen test (CLEIA/ECLEIA)  
抗原定量検査

× **Antigen (test/kit)**  
(抗原検査)

× **Rapid antigen (test/kit)**  
(迅速抗原検査)

× **Antibody (test/kit)**  
(抗体検査)

※日本で無症状者への検査として推奨している検体・検査方法。

※日本で無症状者への検査として推奨されていない検体・検査方法。

### 検査時間

○ **検体採取**が  
出国前の72時間以内

× **結果判明**が  
出国前の72時間以内

※今後、国内外の状況に鑑み、上記取扱いを変更する可能性があります。

参考：新型コロナウイルス感染症（COVID-19）病原体検査の指針第4版 (<https://www.mhlw.go.jp/content/000788513.pdf>)



# Requirements for Certificate of Testing for Entering Japan

※Only test certificates which show the results of the test using valid samples and testing methods below are valid.

## Valid

### Sample:

- Nasopharyngeal/Nasopharynx (Swab/smear)
- **Rhinopharyngeal/Rhinopharynx** (Swab/smear)
- (Deep throat)Saliva sample
- **Nasopharyngeal (※) and oropharyngeal(throat) (swab /smear)**  
(Naso and oropharyngeal/Rhino and oropharyngeal/oro and nashopharyngeal (※) )  
(※Nasopharyngeal/Rhinopharyngeal/Nasopharynx/Rhinopharynx)

### Testing Method:

#### Nucleic acid amplification test

- PCR - Polymerase chain reaction
- (RT-)PCR - Reverse transcription PCR  
(**real-time (RT-)PCR/Q-PCR/Fluorescence-PCR/Multiplex-PCR**)
- LAMP – Loop-mediated isothermal amplification (**RT-LAMP**)
- TMA – Transcription mediated amplification
- TRC - Transcription-reverse transcription concerted reaction
- Smart Amp - Smart amplification process
- NEAR - Nicking endonuclease amplification reaction (**e.g. ID-NOW**)

#### Others

- Next generation sequencing
- ~~Quantitative antigen test (CLEIA/**ECLEIA**)~~

※Recommend in Japan as a sample/testing method for asymptomatic people.

## Invalid

- Nasal (swab/smear)
  - Oral (swab/smear)
  - Throat (swab/smear)
  - Mid-Turbinate Nasal (swab/smear)
  - Nose
  - Gargle Water
  - Mixture of sample “A” and “B”  
“A” and “B” show all types of sample(**except nasopharyngeal and oropharyngeal(swab/smear)**).
- <Example>
- Nasal and throat (swab/smear)
  - Pharyngeal and nasal (swab/smear)
  - Nasal and oropharyngeal/oropharynx (swab/smear)

- Antigen test/test kit
- Rapid antigen test/test kit
- Antibody test/test kit

※Not recommend in Japan as a sample/testing method for asymptomatic people.

**Sample collection** must be done Within 72 hours before the departure time of the flight.

※These requirements may be changed depending on infection status.